

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

101734-CN					
ID	Weight %	Concentration (mg/g)			
<b>∆9-THC</b>	0.0331	0.331			
THCV	ND	ND			
CBD	1.58	15.8			
CBDV	0.0122	0.122			
CBG	0.0178	0.178			
CBC	0.0579	0.579			
CBN	<loq< td=""><td><loq< td=""><td></td><td></td><td></td></loq<></td></loq<>	<loq< td=""><td></td><td></td><td></td></loq<>			
THCA	0.0109	0.109			
CBDA	0.417	4.17			
CBGA	0.0111	0.111			
∆8-THC	ND	ND			
exo-THC	ND	ND			
Total	2.14	21.4	0%	Cannabinoids (wt%)	1.58%
Max THC	0.0427	0.427		Limit of Quantitation $(LOQ) = 0$	0.0100 wt%
Max CBD	1.95	19.5		Limit of Detection $(LOD) = 0$	0.0033 wt%
Ratio of Total (	<b>CBD to THC</b> 45.6:1				

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation:  $MAX THC = (0.877 \times THCA) + THC$ . This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND=None detected above the limits of detection (LOD), which is one third of Limit of Quantification (LOQ). For values reported as "<LOQ", the estimated value is included in the calculated Total.

## **END OF REPORT**

420 Fortune Blvd • Milford, MA 01757 • 617-221-3356 www.ProVerdeLabs.com